arXiv:1602.08667v4 [math.GR] 3 Nov 2016

PROOF OF SOME PROPERTIES OF TRANSFER USING NONCOMMUTATIVE DETERMINANTS

NAOYA YAMAGUCHI

ABSTRACT. A transfer is a group homomorphism from a finite group to an abelian quotient group of a subgroup of the group.

In this paper, we explain some of the properties of transfers by using noncommutative determinants. These properties enable us to understand transfers more naturally.

1. INTRODUCTION

A transfer is defined by Issai Schur [6] as a group homomorphism from a finite group to the abelianization of a subgroup of the group. In finite group theory, transfers play an important role in transfer theorems. Transfer theorems include, for example, Alperin's theorem [1, Theorem 4.2], Burnside's theorem [5, Hauptsatz], and Hall-Wielandt's theorem [4, Theorem 14.4.2].

On the other hand, Eduard Study defined the determinant of a quaternionic matrix [3]. The Study determinant uses a regular representation from $\mathrm{Mat}(n, \mathbb{H})$ to $\mathrm{Mat}(2n, \mathbb{C})$, where $\mathbb{H}$ is the set of quaternions. Similarly, we define a noncommutative determinant. It is similar to the Dieudonné determinant [2].

Tôru Umeda suggested that a transfer can be derived as a noncommutatvie determinant [7, Footnote 7]. In this paper, we develop his ideas in order to explain the properties of the transfers by using noncommutative determinants. As a result, we can understand the transfers more naturally.

Let G be a group, H a subgroup of G of finite index, K a normal subgroup of H, and the quotient group H/K of K in H an abelian group. The transfer of G into H/K is a group homomorphism $V_{G\to H/K} : G \to H/K$. The definition of the transfer $V_{G\to H/K}$ uses the left (or right) coset representatives of H in G. We can show that a transfer has the following properties.

(1) A transfer is a group homomorphism from G to H/K.
(2) A transfer is an invariant under change of coset representatives.
(3) A transfer by left coset representatives equals a transfer by right coset representatives.

Let R be a commutative ring with unity and RG the group algebra of G over R whose elements are all possible finite sums of the form $\sum_{g\in G} x_g g, x_g \in R$. The noncommutative determinant uses a left (or right) regular representation from RG to $\mathrm{Mat}(m, RH)$, where m is the index of H in G. We can show that the noncommutative determinant has the following properties.

Date: November 4, 2016.
2010 *Mathematics Subject Classification.* Primary 20C05; Secondary 11H60; 15A15.
Key words and phrases. transfer; noncommutative determinant; group algebra.

($1'$) The determinant is a homomorphism from RG to H/K.
($2'$) The determinant is an invariant under a change of a regular representation.
($3'$) Any left regular representation is equivalent to any right regular representation.

Here, our objective is to obtain the properties of transfers (1), (2), and (3) by using the properties of noncommutative determinants ($1'$), ($2'$), and ($3'$).

2. Definition of the transfer

Here, we define the left and right transfer of G into H/K.

Let $G = t_1H \cup t_2H \cup \cdots \cup t_mH$. That is, we take a complete set $\{t_1, t_2, \ldots, t_m\}$ of left coset representatives of H in G.

We define $\overline{g} = t_i$ for all $g \in t_iH$.

Definition 1 (Left transfer)**.** *We define the map $V_{G\to H} : G \to H/K$ by*

$$V_{G\to H/K}(g) = \prod_{i=1}^{m} \left\{ \left(\overline{gt_i}\right)^{-1} gt_i \right\} K.$$

We call the map $V_{G\to H/K}$ the left transfer of G into H/K.

Next, we define the right transfer of G into H/K.

Let $G = Hu_1 \cup Hu_2 \cup \cdots \cup Hu_m$. That is, we take a complete set $\{u_1, u_2, \ldots, u_m\}$ of right coset representatives of H in G.

We define $\widetilde{g} = u_i$ for all $g \in Hu_i$.

Definition 2 (Right transfer)**.** *We define the map $V_{G\to H/K} : G \to H/K$ by*

$$V_{G\to H/K}(g) = \prod_{i=1}^{m} \left\{ u_i g \left(\widetilde{u_i g}\right)^{-1} \right\} K.$$

We call the map $V_{G\to H/K}$ the right transfer of G into H/K.

The definitions of the left and right transfers use the coset representatives of H in G. But, we can show that the left and right transfers are invariant under a change of coset representatives. Furthermore, we can show that a transfer is a group homomorphism from G to K and a transfer by left coset representatives equals the transfer by right coset representatives.

3. Definition of the noncommutative determinant

Here, we define the noncommutative determinant.

First, we define the left regular representation of RG. We take a complete set $T = \{t_1, t_2, \ldots, t_m\}$ of left coset representatives of H in G.

Definition 3 (Left regular representation)**.** *For all $\alpha \in RG$, there exists a unique $L_T(\alpha) \in \mathrm{Mat}(m, RH)$ such that*

$$\alpha(t_1\ t_2\ \cdots\ t_m) = (t_1\ t_2\ \cdots\ t_m)L_T(\alpha).$$

We call the map $L_T : RG \ni \alpha \mapsto L_T(\alpha) \in \mathrm{Mat}(m, RH)$ the left regular representation with respect to T.

Obviously, L_T is an R-algebra homomorphism.

Let $T' = \{t'_1, t'_2, \ldots, t'_m\}$ be an another complete set of left coset representatives of H in G. Then, there exists $P \in \mathrm{Mat}(m, RH)$ such that $L_T = P^{-1}L_{T'}P$.

Example 4. *Let $G = \mathbb{Z}/2\mathbb{Z} = \{0, 1\}$, $H = \{0\}$ and $\alpha = x0 + y1 \in RG$. Then, we have*

$$(x0 + y1)(0\ 1) = (0\ 1) \begin{bmatrix} x0 & y0 \\ y0 & x0 \end{bmatrix}.$$

To get an expression for L_T, we define the map $\dot{\chi}$ by

$$\dot{\chi}(g) = \begin{cases} 1 & g \in H, \\ 0 & g \notin H \end{cases}$$

for all $g \in G$.

Lemma 5. *Let $\alpha = \sum_{g \in G} x_g g$. Then, we have*

$$L_T(\alpha)_{ij} = \sum_{g \in G} \dot{\chi}\left(t_i^{-1} g t_j\right) x_g t_i^{-1} g t_j.$$

Proof. Furthermore, we have

$$\begin{aligned}
&(t_1\ t_2\ \cdots\ t_m) \left(\sum_{g \in G} \dot{\chi}\left(t_i^{-1} g t_j\right) x_g t_i^{-1} g t_j \right)_{1 \le i \le m, 1 \le j \le m} \\
&= \left(\sum_{i=1}^{m} \sum_{g \in G} \dot{\chi}(t_i^{-1} g t_1) x_g g t_1 \ \sum_{i=1}^{m} \sum_{g \in G} \dot{\chi}(t_i^{-1} g t_2) x_g g t_2 \ \cdots \ \sum_{i=1}^{m} \sum_{g \in G} \dot{\chi}(t_i^{-1} g t_m) x_g g t_m \right) \\
&= \left(\sum_{g \in G} x_g g \right) (t_1\ t_2\ \cdots\ t_m).
\end{aligned}$$

This completes the proof. □

From Lemma 5, we have

$$\begin{aligned}
L_T(g)_{ij} &= \dot{\chi}\left(t_i^{-1} g t_j\right) t_i^{-1} g t_j \\
&= \begin{cases} t_i^{-1} g t_j & t_i^{-1} g t_j \in H, \\ 0 & t_i^{-1} g t_j \notin H. \end{cases}
\end{aligned}$$

From $t_i^{-1} g t_j \in H$ if and only if $\overline{g t_j} = t_i$, we have

$$L_T(g)_{ij} = \begin{cases} \left(\overline{g t_j}\right)^{-1} g t_j & t_i^{-1} g t_j \in H, \\ 0 & t_i^{-1} g t_j \notin H. \end{cases}$$

As for the definition of the noncommutative determinant, let $\psi : \mathrm{Mat}(m, RH) \to \mathrm{Mat}\left(m, R\left(H/K\right)\right)$ be a map such that

$$\psi\left((x_{ij})\right)_{1 \le i \le m, 1 \le j \le m} = \left(x_{ij} K\right)_{1 \le i \le m, 1 \le j \le m}.$$

Obviously, ψ is an R-algebra homomorphism.

Definition 6. *We define the map* $\mathrm{Det} : \mathrm{Mat}(m, RG) \to R\left(H/K\right)$ *by*

$$\mathrm{Det} = \det \circ \psi \circ L_T.$$

Since there is P such that $L_T = P^{-1}L_{T'}P$, we have

$$\begin{aligned}\mathrm{Det} &= \det \circ L_T \\ &= \det \circ L_{T'}.\end{aligned}$$

Thus, the determinant is an invariant under a change of left regular representations, so the determinant Det is well-defined. If K is the commutator subgroup of H, the determinant is similar to the Dieudonné determinant.

Obviously, the map Det is a homomorphism. That is, $\mathrm{Det}(\alpha\beta) = \mathrm{Det}(\alpha)\mathrm{Det}(\beta)$ for all $\alpha, \beta \in RG$. Therefore, we obtain properties $(1')$ and $(2')$.

Remark 7. *In general, that $\alpha \in RG$ is invertible is not equivalent to that $\mathrm{Det}(\alpha) \in R(H/K)$ is invertible. For example, let $R = \mathbb{C}$, $\mathbb{Z}/2\mathbb{Z} = \{\overline{0}, \overline{1}\}$, dihedral group $D_3 = \langle a, b \mid a^3 = b^2 = e, ab = ba^{-1}\rangle$ where e is the unit element of D_3, $G = \mathbb{Z}/2\mathbb{Z} \times D_3$, $H = D_3$ and $K = [D_3, D_3]$ the commutator subgroup of H. Then $\alpha = (\overline{0}, e) + (\overline{0}, a) + (\overline{0}, a^2)$ is not invertible. But, $\mathrm{Det}(\alpha) = 9K$ is invertible.*

4. Proof of the properties

Here, we prove the transfer properties by using the noncommutative determinant's properties.

For all $g \in G$ and for all $t \in T$, there exists a unique $t_j \in T$ such that $t_i^{-1}gt_j \in H$. Therefore, there exists $\mathrm{sgn}(g) \in \{-1, 1\}$ such that

$$\begin{aligned}\mathrm{Det}(g) &= \det\left(\psi\left(L_T(g)\right)\right) \\ &= \mathrm{sgn}(g)\prod_{i=1}^{m}\left\{\left(\overline{gt_i}\right)^{-1}gt_i\right\}K \\ &= \mathrm{sgn}(g)V_{G\to H/K}(g).\end{aligned}$$

Thus, we have

$$\begin{aligned}\mathrm{sgn}(gh)V_{G\to H/K}(gh) &= \mathrm{Det}(gh) \\ &= \mathrm{Det}(g)\mathrm{Det}(h) \\ &= \mathrm{sgn}(g)\mathrm{sgn}(h)V_{G\to H/K}(g)V_{G\to H/K}(h).\end{aligned}$$

Hence, we obtain

$$\begin{aligned}\mathrm{sgn}(gh) &= \mathrm{sgn}(g)\mathrm{sgn}(h), \\ V_{G\to H/K}(gh) &= V_{G\to H/K}(g)V_{G\to H/K}(h).\end{aligned}$$

Therefore, from property $(1')$ that Det is a homomorphism, the left transfer $V_{G\to H/K}$ is a group homomorphism (Assuming, that is, $R = \mathbb{F}_2$, and we do not consider the signature).

Next, we show that the left transfer is an invariant under a change of coset representatives by using property $(2')$ that the determinant is an invariant under a change of regular representations. That is, we show that

$$\prod_{i=1}^{m}\left\{\left(\overline{gt_i}\right)^{-1}gt_i\right\}K = \prod_{i=1}^{m}\left\{\left(\overline{\overline{gt_i'}}\right)^{-1}gt_i'\right\}K$$

where we define $\overline{\overline{g}} = t_i'$ for all $g \in t_i'H$.

From property $(2')$, there exists $\mathrm{sgn}'(g) \in \{-1, 1\}$ such that

$$\begin{aligned}\prod_{i=1}^{m} \left\{ \left(\overline{gt_i}\right)^{-1} gt_i \right\} K &= \mathrm{sgn}(g)\mathrm{Det}(g) \\ &= \mathrm{sgn}(g)\mathrm{sgn}'(g) \prod_{i=1}^{m} \left\{ \left(\overline{\overline{gt'_i}}\right)^{-1} gt'_i \right\} K.\end{aligned}$$

Therefore, we have $\mathrm{sgn}(g)\mathrm{sgn}'(g) = 1$ and

$$\prod_{i=1}^{m} \left\{ \left(\overline{gt_i}\right)^{-1} gt_i \right\} K = \prod_{i=1}^{m} \left\{ \left(\overline{\overline{gt'_i}}\right)^{-1} gt'_i \right\} K.$$

Hence, the left transfer is an invariant under a change of coset representatives.

Now let us prove property (3) that $V_{G\to H/K} = \widetilde{V}_{G\to H/K}$ from property $(3')$ that any left regular representation is equivalent to any right regular representation.

Let $G = Hu_1 \cup Hu_2 \cup \cdots \cup Hu_m$. That is, we take a complete set $U = \{u_1, u_2, \ldots, u_m\}$ of right coset representatives of H in G.

Definition 8. *For all $\alpha \in RG$, there exists $R_U(\alpha) \in \mathrm{Mat}(m, RH)$ such that*

$$\begin{pmatrix} u_1 \\ u_2 \\ \vdots \\ u_m \end{pmatrix} \alpha = R_U(\alpha) \begin{pmatrix} u_1 \\ u_2 \\ \vdots \\ u_m \end{pmatrix}.$$

We call the map $R_U : RG \ni \alpha \mapsto R_U(\alpha) \in \mathrm{Mat}(m, RH)$ the right regular representation.

The same as the left transfer, we can show that the following lemma.

Lemma 9. *Let $\alpha = \sum_{g\in G} x_g g$. Then we have*

$$R_U(\alpha)_{ij} = \sum_{g\in G} \dot{\chi}(u_i g u_j^{-1}) x_g u_i g u_j^{-1}.$$

Therefore, there exists $\widetilde{\mathrm{sgn}}(g) \in \{-1, 1\}$ such that

$$\left(\det \circ \psi \circ R_U\right)(g) = \widetilde{\mathrm{sgn}}(g) \widetilde{V}_{G\to H/K}(g)$$

and $\widetilde{V}_{G\to H/K}$ is an invariant under a change of coset representatives of H in G. We have properties (1) and (2).

Since T is a complete set of left coset representatives of H in G, we can take a complete set of $T^{-1} = \{t_1^{-1}, t_2^{-1}, \ldots, t_m^{-1}\}$ of right coset representatives of H in G. Therefore,

$$\begin{aligned} R_{T^{-1}}(\alpha)_{ij} &= \sum_{g\in G} \dot{\chi}\left(t_i^{-1} g (t_j^{-1})^{-1}\right) x_g t_i^{-1} g (t_j^{-1})^{-1} \\ &= L_T(\alpha)_{ij}. \end{aligned}$$

We obtain property $(3')$. As a result,

$$\left(\det \circ \psi \circ R_U\right)(g) = \left(\det \circ \psi \circ L_T\right)(g).$$

Therefore, we have

$$\widetilde{\mathrm{sgn}}(g) = \mathrm{sgn}(g),$$
$$\widetilde{V}_{G\to H/K} = V_{G\to H/K}.$$

We obtain property (3).

Acknowledgments I am deeply grateful to Prof. Tôru Umeda, Prof. Hiroyuki Ochiai and Prof. Hideaki Morita who provided helpful comments and suggestions. In particular, Tôru Umeda gave me the initial motivation for undertaking this study. I would also like to thank my colleagues in the Graduate School of Mathematics of Kyushu University, in particular Cid Reyes for comments and suggestions. This work was supported by a grant from the Japan Society for the Promotion of Science (JSPS KAKENHI Grant Number 15J06842).

REFERENCES

[1] ALPERIN, Jonathan L. Sylow intersections and fusion. *Journal of Algebra*, 1967, 6.2: 222–241.
[2] ARTIN, Emil. *Geometric algebra*. Courier Dover Publications, 2016.
[3] ASLAKSEN, Helmer. Quaternionic determinants. *The Mathematical Intelligencer*, 1996, 18.3: 57–65.
[4] HALL, Marshall. *The theory of groups*. American Mathematical Soc., 1976.
[5] HUPPERT, B. Endliche gruppen I. *Spring-Verlag*, New York Berlin, 1967.
[6] SCHUR, I. Neuer Beweis eines Satzes über endliche Gruppen, *Sitzungsberichteder Königlich Preussischen Akademie der Wissenschaften zu Berlin*, 1902, 1013–1019.
[7] UMEDA, Tôru, On some variants of induced representations, Symposium on Representation Theory 2012, 2012, 7–17 (in Japanese).
[8] YOSHIDA, Tomoyuki. Character-theoretic transfer. *Journal of Algebra*, 1978, 52.1: 1–38.

Naoya Yamaguchi
Graduate School of Mathematics
Kyushu University
Nishi-ku, Fukuoka 819-0395
Japan
n-yamaguchi@math.kyushu-u.ac.jp